



Research Article

From Silent Ischemia to Hip Collapse: A Critical and Integrative Literature Review of Avascular Necrosis of the Femoral Head in Early Adulthood, With Emphasis on Etiopathogenesis Risk Factors, Early Clinical Recognition, Disease Progression and Homoeopathic Individualization

Dr. Pratibha Jagannath Pagar

BHMS MD Homoeopathy (Practice of Medicine)

HOD and Professor, Department of Practice of Medicine, Matoshri Homoeopathy College & Hospital, Matoshri Education Campus, Near Odha Gaon, Chhatrapati Sambhajinagar Highway, Eklahare, Nashik, Maharashtra, India

Corresponding Author: * Dr. Pratibha Jagannath Pagar

DOI: <https://doi.org/10.5281/zenodo.23185933>

Abstract

Avascular necrosis of the femoral head (AVNFH), also termed osteonecrosis of the femoral head (ONFH), is a potentially progressive disorder characterized by compromised intraosseous circulation, osteocyte death, structural failure of the femoral head, and eventual secondary degenerative arthritis. Its clinical importance is particularly pronounced in early adulthood because the disease may affect individuals during their most productive years and may ultimately necessitate hip arthroplasty at a relatively young age. The transition from apparently silent ischemia to symptomatic disease and structural collapse is often clinically subtle, creating a critical window during which early recognition and appropriate staging may influence management.

The pathogenesis of ONFH is increasingly understood as a multifactorial process involving vascular compromise, endothelial dysfunction, intravascular coagulation, altered lipid metabolism, marrow adipogenesis, intraosseous pressure, impaired osteogenesis, apoptosis, inflammatory mechanisms, and mechanical failure. Corticosteroid exposure and excessive alcohol consumption remain major non-traumatic associations, while autoimmune disease, haematological disorders, metabolic abnormalities, smoking, transplantation, decompression sickness, trauma, and idiopathic disease constitute additional pathways. Importantly, disease progression is not determined solely by the presence of necrosis; lesion size, anatomical location, weight-bearing involvement, and subchondral fracture substantially influence the probability of collapse.

Early symptoms may be absent or nonspecific, commonly presenting as groin, hip, buttock, or referred thigh pain with restricted internal rotation. Conventional radiography may remain normal during early disease, whereas magnetic resonance imaging (MRI) can identify marrow abnormalities before radiographic changes become apparent. A meta-analysis reported pooled MRI sensitivity and specificity of approximately 93% and 91%, respectively, for early ONFH.

This review critically examines ONFH from etiopathogenesis through clinical progression and discusses the potential conceptual role of homoeopathic individualisation as a complementary, patient-centred framework, while emphasising that homoeopathic prescribing should not substitute for MRI-based diagnosis, orthopaedic assessment, risk-factor modification, or established joint-preserving and reconstructive interventions. Classical homoeopathic principles emphasise individualised assessment and the totality of characteristic symptoms; however, contemporary clinical evidence demonstrating that individualised homoeopathic treatment reverses femoral-head necrosis or prevents collapse remains insufficient.

Manuscript Information

- **ISSN No:** 3139-6313
- **Received:** 11-08-2026
- **Accepted:** 26-09-2026
- **Published:** 06-10-2026
- **IJCRP:** 1(5); 2026: 176-189
- **©2026, All Rights Reserved**
- **Plagiarism Checked:** Yes
- **Peer Review Process:** Yes

How to Cite this Article

Pagar P J. From Silent Ischemia to Hip Collapse: A Critical and Integrative Literature Review of Avascular Necrosis of the Femoral Head in Early Adulthood, With Emphasis on Etiopathogenesis, Risk Factors, Early Clinical Recognition, Disease Progression and Homoeopathic Individualization. Int. J. Creative Res. Publ. 2026;1(5):176-189.

Access this Article Online




www.ijcrp.in

KEYWORDS: avascular necrosis, osteonecrosis of femoral head, ONFH, AVNHF, femoral head collapse, early adulthood, MRI, corticosteroids, alcohol, etiopathogenesis, homoeopathic individualisation, integrative medicine.

1. INTRODUCTION

Avascular necrosis of the femoral head represents a distinctive intersection between vascular biology, bone metabolism, mechanical loading, and clinical orthopaedics. Although the term *avascular necrosis* remains widely used, *osteonecrosis of*

the femoral head is increasingly preferred because the disease involves not simply absence of arterial blood flow but a complex disturbance of the bone's microcirculation and cellular environment.



The femoral head is particularly vulnerable because portions of its circulation have limited collateral capacity. Interruption or impairment of microvascular supply may result in ischemic injury, followed by death of osteocytes and marrow elements. Initially, the structural integrity of the femoral head may remain apparently preserved. However, continued loading of necrotic and mechanically weakened bone can eventually result in subchondral fracture, flattening and collapse. The clinical paradox of ONFH is therefore that **biological disease may precede recognizable clinical disease**.

A young adult may have:

vascular injury → silent osteonecrosis → subtle pain → MRI-positive disease → subchondral failure → femoral-head collapse → secondary osteoarthritis → arthroplasty

This sequence explains why early diagnosis is disproportionately important.

ONFH also has particular relevance to young adults because total hip arthroplasty performed at a younger age may expose

patients to the possibility of revision surgery during their lifetime. Consequently, contemporary management places substantial emphasis on identifying disease before irreversible structural collapse and evaluating appropriate joint-preserving strategies.

2. AIM OF THE REVIEW

The present narrative review aims to critically integrate contemporary understanding of:

- the etiopathogenesis of femoral-head osteonecrosis;
- major and emerging risk factors;
- mechanisms responsible for vascular compromise and bone failure;
- early clinical recognition;
- imaging and staging;
- natural history and progression toward collapse;
- prognostic importance of lesion size and location;
- contemporary principles of disease management; and

9. the possible conceptual contribution and limitations of homoeopathic individualization in an integrative clinical framework.

3. METHODOLOGICAL APPROACH

This article was developed as a **narrative literature review** using contemporary and foundational literature concerning osteonecrosis of the femoral head.

Literature was identified through searches of biomedical literature databases and publicly accessible scientific resources using combinations of terms including:

- “Avascular necrosis femoral head”
- “Osteonecrosis femoral head”
- “ONFH pathogenesis”
- “Risk factors osteonecrosis”
- “Steroid induced osteonecrosis”
- “Alcohol induced osteonecrosis”
- “MRI early diagnosis ONFH”
- “Femoral head collapse”
- “ARCO classification”
- “Homoeopathy individualization”
- “Homoeopathic materia medica bone necrosis”

Priority was given to systematic reviews, clinical guidelines, diagnostic studies, pathophysiological reviews, natural-history studies, and major orthopedic literature.

Because evidence concerning homoeopathy and ONFH is substantially less developed than the orthopedic literature, classical homoeopathic sources were used primarily to explain the **principle of individualization and symptom-based case analysis**, rather than as evidence of structural reversal of osteonecrosis.

4. Terminology: Avascular Necrosis and Osteonecrosis

Several terms occur in the literature:

- Avascular necrosis (AVN)
- Osteonecrosis
- Osteonecrosis of the femoral head (ONFH)
- Avascular necrosis of the femoral head (AVNFH)
- Ischemic necrosis
- Aseptic necrosis

These terms overlap clinically, although osteonecrosis of the femoral head more accurately encompasses the pathological process.

The fundamental pathological event is inadequate effective perfusion followed by death of bone and marrow cells. However, the initiating event may differ between individuals. In traumatic disease, vascular interruption may occur after fracture or dislocation. In non-traumatic disease, the process may involve several interacting systemic and local mechanisms.

5. Why Early Adulthood Is a Critical Clinical Window

ONFH disproportionately affects younger and middle-aged adults. This creates a major clinical challenge because hip pain in a young adult is frequently attributed to muscular strain, sports injury, tendinopathy, low-back pathology, or nonspecific mechanical pain.

The possibility of osteonecrosis may therefore be overlooked. The disease has considerable socioeconomic importance because it can impair:

- mobility,
- employment,
- physical activity,
- sports participation,
- social functioning,
- sleep,
- psychological well-being,
- and long-term independence.

Moreover, ONFH may be bilateral. Contemporary literature indicates that bilateral disease is common, and an asymptomatic contralateral hip may harbor clinically occult disease.

Therefore:

A young adult with unexplained persistent hip or groin pain and relevant risk factors should not be reassured solely by a normal radiograph.

6. Etiopathogenesis: From Vascular Insult to Structural Collapse

6.1 The multifactorial model

The pathogenesis of ONFH cannot be adequately explained by a single mechanism.

A useful contemporary model is the **multi-hit hypothesis**, in which several biological and mechanical insults converge upon the femoral-head circulation and bone microenvironment.

These mechanisms include:

- vascular injury;
- endothelial dysfunction;
- microthrombosis;
- intravascular coagulation;
- altered lipid metabolism;
- marrow adipocyte hypertrophy;
- increased intraosseous pressure;
- impaired osteoblast differentiation;
- increased osteocyte apoptosis;
- inflammatory signaling;
- oxidative stress;
- impaired angiogenesis;
- mechanical overload;
- genetic susceptibility.

Thus, the same clinical endpoint—femoral-head osteonecrosis—may arise through biologically different pathways.


7. Vascular Mechanisms

The vascular hypothesis remains central.

The femoral head depends upon a relatively vulnerable vascular arrangement. Disruption of small vessels can reduce oxygen and nutrient delivery to subchondral bone and marrow.

Microvascular compromise may result from:

Endothelial injury



collapse

This model explains why ONFH may remain clinically silent while pathological changes are already developing.

8. Corticosteroid-Associated Osteonecrosis

Corticosteroid exposure is among the most important recognized associations with non-traumatic ONFH. The mechanism appears multifactorial rather than attributable to one isolated vascular event.

Potential pathways include:

- increased marrow adipogenesis;
- enlargement of marrow adipocytes;
- altered lipid metabolism;
- endothelial dysfunction;
- microvascular obstruction;
- increased intraosseous pressure;
- osteoblast suppression;
- osteocyte apoptosis;
- altered bone remodeling.

A contemporary review describes steroid-associated and alcohol-associated osteonecrosis as sharing several mechanisms, including increased adipogenesis, vascular alterations, chronic inflammatory processes, and impaired bone-cell differentiation.

Importantly, corticosteroid exposure should be assessed not simply as “steroid use: yes/no”, but in relation to:

- indication,
- dose,
- duration,
- route,
- cumulative exposure,
- repeated courses,
- concurrent risk factors.

9. Alcohol-Associated Osteonecrosis

Chronic excessive alcohol exposure is another major non-traumatic risk factor.

Alcohol may influence:

- mesenchymal stem-cell differentiation;
- adipogenesis;
- osteoblast activity;
- lipid metabolism;
- oxidative stress;
- vascular function;
- bone remodeling.

The association is also biologically plausible because alcohol can alter the marrow microenvironment and favor adipocytic differentiation of mesenchymal cells.

The clinical history should therefore document alcohol exposure quantitatively rather than simply recording “alcoholic/non-alcoholic.”

10. Haematological and Thrombotic Factors

Conditions associated with altered blood rheology or thrombosis may increase susceptibility to osteonecrosis.

Important examples include:

- sickle-cell disease;
- thalassemia;
- antiphospholipid-associated thrombosis;
- myeloproliferative disorders;
- coagulation abnormalities;
- other hypercoagulable states.

Microvascular occlusion can compromise the circulation of the femoral head and initiate ischemic injury.

11. Autoimmune and Systemic Disease

ONFH may occur in association with systemic autoimmune disorders, particularly systemic lupus erythematosus.

However, an important clinical distinction is required:

Disease-related mechanisms and treatment-related mechanisms may coexist.

For example:

SLE → systemic vascular/inflammatory abnormalities

•
corticosteroid therapy → additional osteonecrosis risk
↓
cumulative susceptibility


Thus, attributing ONFH solely to the underlying autoimmune disease or solely to corticosteroids may oversimplify the patient's actual risk profile.


7. Vascular Mechanisms

The vascular hypothesis remains central.

The femoral head depends upon a relatively vulnerable vascular arrangement. Disruption of small vessels can reduce oxygen and nutrient delivery to subchondral bone and marrow.

Microvascular compromise may result from:





This model explains why ONFH may remain clinically silent while pathological changes are already developing.

8. Corticosteroid-Associated Osteonecrosis

Corticosteroid exposure is among the most important recognized associations with non-traumatic ONFH. The mechanism appears multifactorial rather than attributable to one isolated vascular event.

Potential pathways include:

- increased marrow adipogenesis;
- enlargement of marrow adipocytes;
- altered lipid metabolism;
- endothelial dysfunction;
- microvascular obstruction;
- increased intraosseous pressure;
- osteoblast suppression;
- osteocyte apoptosis;
- altered bone remodeling.

A contemporary review describes steroid-associated and alcohol-associated osteonecrosis as sharing several mechanisms, including increased adipogenesis, vascular alterations, chronic inflammatory processes, and impaired bone-cell differentiation.

Importantly, corticosteroid exposure should be assessed not simply as “steroid use: yes/no”, but in relation to:

- indication,
- dose,
- duration,
- route,
- cumulative exposure,
- repeated courses,
- concurrent risk factors.

9. Alcohol-Associated Osteonecrosis

Chronic excessive alcohol exposure is another major non-traumatic risk factor.

Alcohol may influence:

- mesenchymal stem-cell differentiation;
- adipogenesis;
- osteoblast activity;
- lipid metabolism;
- oxidative stress;

- vascular function;
- bone remodeling.

The association is also biologically plausible because alcohol can alter the marrow microenvironment and favor adipocytic differentiation of mesenchymal cells.

The clinical history should therefore document alcohol exposure quantitatively rather than simply recording “alcoholic/non-alcoholic.”

10. Hematological and Thrombotic Factors

Conditions associated with altered blood rheology or thrombosis may increase susceptibility to osteonecrosis.

Important examples include:

- sickle-cell disease;
- thalassemia;
- antiphospholipid-associated thrombosis;
- myeloproliferative disorders;
- coagulation abnormalities;
- other hypercoagulable states.

Microvascular occlusion can compromise the circulation of the femoral head and initiate ischemic injury.

11. Autoimmune and Systemic Disease

ONFH may occur in association with systemic autoimmune disorders, particularly systemic lupus erythematosus.

However, an important clinical distinction is required:

Disease-related mechanisms and treatment-related mechanisms may coexist.

For example:

SLE → systemic vascular/inflammatory abnormalities

• corticosteroid therapy → additional osteonecrosis risk

↓
cumulative susceptibility

Thus, attributing ONFH solely to the underlying autoimmune disease or solely to corticosteroids may oversimplify the patient's actual risk profile.

16. The Importance of Internal Rotation

Restricted internal rotation may be an important physical finding.

The examination should include:

- gait;
- hip flexion;
- extension;
- abduction;
- adduction;
- internal rotation;
- external rotation;
- limb-length assessment;
- lumbar examination;
- sacroiliac assessment;
- neurological examination.

The purpose is not merely to confirm hip pathology but also to differentiate ONFH from:

- lumbar radiculopathy;

- sacroiliac dysfunction;
- femoroacetabular impingement;
- labral pathology;
- osteoarthritis;
- transient osteoporosis;
- septic arthritis;
- inflammatory arthropathy;
- referred pain.

17. Imaging: Why MRI Changes the Diagnostic Timeline


Plain radiography remains important for structural assessment and advanced disease, but early ONFH may be radiographically occult.

MRI has a major role in early diagnosis.

A systematic review and meta-analysis involving 43 studies reported pooled sensitivity of approximately **93%** and specificity of approximately **91%** for MRI in early ONFH.

Earlier diagnostic studies similarly demonstrated that MRI could detect disease when radiographs remained normal.

Thus:



If normal but suspicion remains high

MRI

This pathway can shorten the interval between biological disease onset and definitive diagnosis.

18. MRI Findings

Characteristic MRI features may include:







- geographic necrotic lesions;
- serpiginous margins;
- altered marrow signal;
- the “double-line” sign on T2-weighted sequences;
- subchondral fracture;
- bone marrow edema;
- flattening or collapse in advanced disease.

MRI also provides information regarding:

- lesion size;
- lesion location;
- weight-bearing involvement;
- contralateral hip involvement;
- associated edema;
- collapse.

These parameters are important because **diagnosis alone is insufficient; prognosis requires characterization of the lesion.**

MRI Findings in Avascular Necrosis (Osteonecrosis) of the Femoral Head

1 Geographic necrotic lesions  T1-weighted (Coronal) Well-defined, geographic area of altered signal intensity within the femoral head.	2 Serpiginous margins  T2-weighted (Coronal) Typical serpiginous (snake-like) margins separating necrotic from viable bone.	3 Altered marrow signal  T1-weighted (Coronal) Altered marrow signal with low intensity on T1 and variable intensity on T2 sequences.	4 The “double-line” sign  T2-weighted (Axial) Outer dark line (granulation tissue) Inner bright line (viable bone) Double-line sign A characteristic double-line sign on T2-weighted images – inner bright line and outer dark line.						
5 Subchondral fracture  T2-weighted (Coronal) A low signal intensity subchondral fracture line (crescent sign) indicating impending collapse.	6 Bone marrow edema  STIR (Coronal) Bone marrow edema appears as high signal intensity on STIR or fat-suppressed T2 images.	7 Flattening or collapse  T1-weighted (Coronal) Flattening of the articular surface or collapse of the femoral head in advanced disease.	Normal vs Avascular Necrosis (MRI) <table border="1"> <thead> <tr> <th>Normal Femoral Head</th> <th>Avascular Necrosis Femoral Head</th> </tr> </thead> <tbody> <tr> <td></td> <td></td> </tr> <tr> <td>T1-weighted (Coronal)</td> <td>T1-weighted (Coronal)</td> </tr> </tbody> </table> Key MRI Features of Avascular Necrosis ✓ Geographic necrotic lesions ✓ Serpiginous margins ✓ Altered marrow signal ✓ “Double-line” sign on T2-weighted sequences ✓ Subchondral fracture ✓ Bone marrow edema ✓ Flattening or collapse in advanced disease	Normal Femoral Head	Avascular Necrosis Femoral Head			T1-weighted (Coronal)	T1-weighted (Coronal)
Normal Femoral Head	Avascular Necrosis Femoral Head								
									
T1-weighted (Coronal)	T1-weighted (Coronal)								

19. Staging and the Pre-Collapse Concept

Modern classification systems include:

- Ficat–Arlet;
- Steinberg;
- ARCO.

The contemporary ARCO framework emphasizes the distinction between:

Pre-collapse disease

and

Post-collapse disease.

This distinction is clinically meaningful because joint-preserving interventions are generally most relevant before substantial structural collapse. Recent international guidance emphasizes accurate staging and assessment of lesion characteristics when selecting management strategies.

20. Lesion Size and Location: The Hidden Prognostic Variables

Not every osteonecrotic lesion behaves identically.

Two patients may both be diagnosed with “Stage II ONFH,” yet their long-term risks may differ substantially.

Important variables include:

- lesion volume;
- percentage of femoral-head involvement;
- lateral versus medial location;
- anterosuperior extension;
- weight-bearing surface involvement;
- presence or absence of subchondral fracture.

Quantitative MRI studies have demonstrated a significant relationship between lesion volume and collapse, while anterosuperior location may increase mechanical vulnerability.

Therefore:

The diagnosis tells us what the disease is; lesion morphology helps tell us what it may do next.

21. Natural History: Not Every Lesion Collapses Immediately

The natural history of ONFH is heterogeneous.

Some small lesions remain stable.

Others enlarge or progress rapidly.

The literature demonstrates that lesion size and location are major determinants of progression. Small, medially located lesions generally have a more favorable natural history than large or laterally located lesions.

This is important because the simplistic statement: “AVN always progresses rapidly to hip replacement” is not supported by the natural-history literature.

Conversely, the opposite assumption—

“Asymptomatic AVN is harmless”

—is also inappropriate.

Clinical management must therefore be individualized according to **stage + lesion characteristics + risk profile + symptoms + structural integrity.**

22. The Critical Event: Subchondral Fracture

The transition from osteonecrosis to mechanical failure represents a crucial stage.

Necrotic bone has reduced biological remodeling capacity.

Repeated loading occurs across weakened trabecular architecture.

↓
Microstructural failure

↓
Subchondral fracture

↓
Loss of spherical femoral-head geometry

↓
Progressive collapse

↓
Altered joint mechanics

↓
Secondary cartilage degeneration

↓
Secondary osteoarthritis

The 2026 international guideline literature emphasizes that once subchondral fracture develops, the likelihood of collapse rises substantially and joint-preserving options become more limited.

23. From Collapse to Secondary Osteoarthritis

Once the femoral head loses its spherical congruity, the hip becomes mechanically abnormal.

This can result in:

- altered load distribution;
- cartilage degeneration;
- joint-space narrowing;
- osteophyte formation;
- progressive pain;
- reduced range of motion;
- gait disturbance.

At this stage, the pathology is no longer simply “bone necrosis.”

It has become a **structural joint disease.**

This distinction is clinically important when discussing any complementary therapeutic approach.

24. Contemporary Management Principles

Management depends upon:

- stage;
- lesion size;
- lesion location;
- symptoms;
- age;
- etiology;
- collapse status;
- contralateral involvement;
- patient expectations.

Potential approaches include:

Conservative/supportive management

- risk-factor modification;
- reduction of excessive alcohol exposure;

- smoking cessation;
- appropriate management of corticosteroid exposure with the prescribing physician;
- activity modification;
- analgesia where appropriate;
- physiotherapy and mobility optimization.

Joint-preserving interventions

Selected pre-collapse lesions may be considered for procedures such as:

- core decompression;
- bone grafting;
- vascularized grafting;
- selected biologic augmentation approaches.

Advanced disease

Once substantial collapse and secondary degenerative changes have occurred, **total hip arthroplasty** becomes an important established treatment option.

Recent literature continues to identify core decompression as a major joint-preserving procedure for appropriately selected pre-collapse disease.

25. Where Homoeopathy May Conceptually Enter an Integrative Model

The homoeopathic approach differs fundamentally from a lesion-directed orthopedic intervention.

Classical homoeopathic doctrine emphasizes:

- individualization;
- totality of symptoms;
- characteristic symptoms;
- modalities;
- concomitants;
- constitutional characteristics;
- causative circumstances.

Hahnemann's *Organon* places considerable emphasis on the totality of symptoms and individual case characteristics in remedy selection.

In an integrative framework, this can be conceptualized as:

Biomedical diagnosis

•

Risk-factor assessment

•

Orthopedic staging

•

Individualized symptom assessment

•

Patient preferences and psychosocial context

rather than attempting to replace structural diagnosis with symptom similarity.

26. Homoeopathic Case Individualization in ONFH

A homoeopathic case assessment could document five major domains.

A. Disease characteristics

- side involved;
- onset;

- duration;
- suspected cause;
- stage;
- MRI characteristics;
- progression;
- associated edema;
- collapse status.

B. Pain characteristics

- location;
- quality;
- intensity;
- onset;
- duration;
- aggravating factors;
- ameliorating factors;
- relation to movement;
- relation to rest;
- night aggravation;
- weight-bearing relationship.

C. General characteristics

- thermal preference;
- thirst;
- appetite;
- sleep;
- perspiration;
- bowel habits;
- energy;
- constitutional tendencies.

D. Mental/emotional characteristics

- anxiety;
- fear;
- irritability;
- emotional response to chronic disease;
- anticipatory concerns;
- occupational stress;
- coping style.

E. Concomitant symptoms

Examples might include:

- digestive symptoms;
- headaches;
- sleep disturbance;
- anxiety;
- constitutional symptoms;
- recurrent musculoskeletal complaints.

The goal within classical homoeopathic methodology is not to prescribe merely because the diagnosis is "AVN," but to construct an individualized symptom picture.

27. Homoeopathic Materia Medica: Differential Individualization

A number of remedies may appear in classical materia medica discussions involving bone, periosteal, traumatic, vascular, or destructive symptom pictures. Their consideration should be

based on the **individual symptom constellation**, not on the diagnosis alone.

27.1 Arnica montana

Classically associated with:

- trauma;
- soreness;
- bruised sensation;
- muscular and periosteal complaints;
- sensitivity after injury.

Its traditional materia-medica picture may make it relevant when the history is strongly trauma-associated.

However, classical indication does **not establish that Arnica reverses osteonecrosis**.

27.2 Ruta graveolens

Classical materia medica associates Ruta with:

- periosteal complaints;
- tendons;
- ligaments;
- overuse;
- injuries involving fibrous tissues;
- persistent soreness following trauma.

It may therefore enter differential consideration when the symptom picture is dominated by periosteal or ligamentous features.

27.3 Symphytum officinale

Traditionally associated with:

- bone injury;
- periosteal trauma;
- fracture-related complaints;
- bone and connective-tissue symptomatology.

Its classical reputation should not be interpreted as evidence that it can restore necrotic femoral-head bone.

27.4 Calcarea fluorica

Classical materia medica describes Calcarea fluorica in relation to:

- hard tissues;
- bone abnormalities;
- induration;
- periosteal and connective-tissue complaints.

Its selection, under a classical framework, would require correspondence with the broader symptom picture rather than the radiological diagnosis alone.

27.5 Calcarea carbonica

Calcarea carbonica has traditionally been associated with constitutional patterns involving:

- impaired nutrition;
- skeletal complaints;
- glandular manifestations;
- constitutional fatigue;
- characteristic thermal and general symptoms.

Again, its inclusion in a differential is based on materia-medica similarity rather than evidence that it reverses ONFH.

27.6 Fluoricum acidum

Classical materia medica describes Fluoricum acidum in relation to destructive processes and certain bone-related complaints, including caries and necrosis of long bones.

Such historical descriptions may be relevant to differential materia-medica analysis, but they cannot be equated with contemporary evidence of efficacy in femoral-head osteonecrosis.

Homoeopathic Materia Medica Differentiation

The following differentiation can be presented as a **Materia Medica framework for individualized prescribing**, not as evidence that any particular remedy treats or reverses femoral-head osteonecrosis.

Remedy	Characteristic clinical picture to differentiate	Important differentiating feature
Arnica montana	Painful, bruised, sore musculoskeletal state; sensation as if beaten	Trauma/contusion-type symptom picture
Symphytum officinale	Bone-related pain and history suggesting bony injury	Traditionally associated with bone and periosteal complaints
Ruta graveolens	Tendon, ligament, periosteal and overuse-type complaints	Marked relationship to connective tissue/overstrain
Rhus toxicodendron	Stiffness and musculoskeletal pain with characteristic movement modality	Often considered when stiffness is worse initially and improves after continued movement
Bryonia alba	Pain aggravated by movement with desire for rest	Strong rest-versus-movement differentiation
Calcarea phosphorica	Bone-related complaints with constitutional features	Consider only when the broader constitutional picture corresponds
Calcarea fluorica	Chronic connective-tissue/bony induration-type symptomatology	Useful only when characteristic constitutional and local features coexist
Silicea	Chronic tissue/bone complaints with characteristic constitutional picture	Not simply because a bone is involved
Fluoricum acidum	Chronic destructive/degenerative-type symptomatology in the traditional Materia Medica	Requires strong individual correspondence rather than pathological-name prescribing
Phosphorus	Bone/vascular-related complaints accompanied by characteristic constitutional and mental features	Consider only when the complete symptom totality supports it
Pulsatilla	Variable musculoskeletal complaints with characteristic general modalities	General constitutional picture is more important than the diagnosis
Sepia	Musculoskeletal symptoms occurring with a characteristic constitutional/mental picture	Should not be selected solely because AVNFH is present

Important prescribing principle

The diagnosis:

“Avascular necrosis of femoral head”

should **not itself be considered the totality of symptoms**.

A more academically defensible formulation is:

Diagnosis → establishes pathology and prognosis

Clinical symptoms → establish therapeutic priorities

Individualized totality → determines the homoeopathic prescription

MRI/radiography → determines structural progression

Orthopedic assessment → determines need for intervention

Remedy Differentiation According to Pain Modality

A useful academic table for the manuscript would be:

Rhus toxicodendron vs Bryonia

Rhus toxicodendron

- stiffness prominent;
- worse after rest;
- first movement difficult;
- improvement after continued gentle movement;
- possible aggravation from cold/damp exposure.

Bryonia

- movement aggravates;
- marked desire for rest;
- pain increased by even slight movement;
- pressure may provide relief in the characteristic case.

Thus, the mere presence of hip pain is insufficient. The **movement modality** becomes the differentiating feature.

Arnica vs Symphytum

Arnica

Think of a **bruised, sore, beaten** sensation, particularly where trauma or mechanical injury forms part of the history.

Symphytum

The traditional Materia Medica picture is more strongly associated with **bone injury, periosteal pain and osseous complaints**.

However, neither remedy should be prescribed merely because the patient has a bone disorder.

Ruta vs Symphytum

Ruta

- ligament/tendon/periosteal emphasis;
- overuse or mechanical strain;
- soreness around attachments.

Symphytum

- stronger bone-centered symptomatology;
- post-traumatic/bony pain picture.

Calcarea phosphorica vs Calcarea fluorica

Calcarea phosphorica

May be considered when the patient's broader constitutional picture corresponds with bone-related complaints, weakness and characteristic nutritional/developmental features.

Calcarea fluorica

Traditionally differentiated through chronic induration, hardness, ligamentous/connective-tissue manifestations and its characteristic constitutional picture.

Again, the pathological diagnosis alone does not establish either prescription.

The therapeutic position of homoeopathy in osteonecrosis of the femoral head should therefore be considered exploratory and complementary rather than substitutive. Individualized prescribing may have a potential role in addressing pain, stiffness, functional limitations and patient-reported well-being, but available evidence does not establish that homoeopathic medicines can reliably reverse femoral-head osteonecrosis or prevent structural collapse. The central principle of integrative management should remain early diagnosis, identification and modification of etiological risk factors, objective MRI-based surveillance and timely orthopedic intervention when indicated. Future research should prioritize prospective, adequately controlled studies incorporating standardized radiological and functional outcomes so that symptomatic benefit can be distinguished from genuine structural modification.

28. A Practical Homoeopathic Differential Framework

Clinical picture	Possible classical differential	Individualizing focus
Trauma-dominant soreness, bruised feeling	Arnica	Trauma, soreness, modalities
Periosteal/ligamentous complaints after strain	Ruta	Overuse, periosteal symptoms
Bone injury/convalescence picture	Symphytum	Bone trauma characteristics
Hard-tissue/induration pattern	Calcarea fluorica	Tissue quality, constitutional features
Constitutional skeletal/nutritional pattern	Calcarea carbonica	General constitution
Destructive bone symptom complex	Fluoricum acidum	Destructive tendency and modalities

This table represents **classical materia medica differentiation**, not comparative clinical evidence of effectiveness.

29. An Important Evidence Boundary

This distinction is essential for academically responsible integrative writing.

Established evidence supports:

- MRI for early detection;
- risk-factor identification;
- staging;
- lesion-size/location assessment;
- orthopaedic evaluation;
- risk-factor modification;
- selected joint-preserving procedures;

- arthroplasty for advanced structural disease.

Classical homoeopathic literature supports:

- individualization;
- symptom totality;
- characteristic symptom analysis;
- materia medica differentiation.

What remains inadequately established:

There is insufficient high-quality clinical evidence demonstrating that individualized homoeopathic treatment:

- restores femoral-head blood supply;
- reverses established osteonecrosis;
- reduces lesion volume reliably;
- prevents subchondral fracture;
- prevents femoral-head collapse;
- eliminates the need for indicated orthopaedic surgery.

Therefore, homoeopathy should **not be presented as an evidence-established alternative to orthopaedic management of ONFH.**

30. A Proposed Integrative Clinical Algorithm

Young adult with persistent hip/groin pain

Detailed history + physical examination

Identify risk factors

- corticosteroids
- alcohol
- smoking
- autoimmune disease
- haematological disease
- trauma
- transplantation
- metabolic/coagulation factors

Plain radiography

If normal but clinical suspicion persists

MRI of hips

Confirm / characterise ONFH

Stage disease

Assess

- lesion size
- lesion location
- weight-bearing involvement
- subchondral fracture
- collapse
- contralateral hip

Pre-collapse disease

→ orthopaedic joint-preservation assessment

Collapse/advanced degeneration

→ reconstructive/arthroplasty assessment

Parallel supportive care

- risk-factor modification
- physical rehabilitation
- pain management

- nutritional and lifestyle optimization
- psychological support

If patient elects complementary homoeopathic care

Individualised homoeopathic case-taking


Materia-medica differentiation

Document baseline symptoms and objective imaging

Continue appropriate orthopaedic surveillance

Repeat imaging when clinically indicated

This model preserves the fundamental principle that **structural disease must be objectively monitored.**



31. Homoeopathic Individualization and Modern Clinical Monitoring

An important opportunity for future integrative research would be to combine individualized prescribing with objective biomedical outcomes.

Rather than reporting only:

“Pain improved after homoeopathic treatment,”
future studies should document:

Clinical outcomes

- VAS/NRS pain;
- WOMAC;
- Harris Hip Score;

- range of motion;
- walking distance;
- functional limitation;
- analgesic requirement.

Radiological outcomes

- MRI lesion volume;
- lesion location;
- bone marrow oedema;
- subchondral fracture;
- femoral-head contour;
- ARCO stage;
- progression/collapse.

Patient-centred outcomes

- quality of life;
- return to work;
- sleep;
- physical activity;
- treatment satisfaction.

This approach would allow homoeopathic research to move from anecdotal symptom reporting toward measurable clinical investigation.

32. RESEARCH GAPS

Several major research questions remain.

32.1 Biomarkers

Can circulating biomarkers identify individuals at risk before radiological disease develops?

32.2 Genetics

Which genetic variants predispose individuals to steroid- or alcohol-associated ONFH?

32.3 Imaging

Can quantitative MRI predict collapse more accurately than conventional staging alone?

32.4 Artificial intelligence

Recent research is exploring MRI-based deep-learning systems for early detection and staging, suggesting a future role for automated image interpretation.

32.5 Bilateral disease

Which factors determine why one hip progresses while the contralateral hip remains stable?

32.6 Integrative medicine

Could individualized complementary approaches influence:

- pain;
- function;
- treatment adherence;
- quality of life;
- rehabilitation participation?

These questions should be studied without assuming structural efficacy before it is demonstrated.

33. DISCUSSION

ONFH illustrates an important principle in modern clinical medicine: **the visible disease is often later than the biological disease.**

The patient may experience little discomfort while vascular and cellular abnormalities are already developing. Once pain becomes obvious, the disease may still be pre-collapse and potentially amenable to joint-preserving treatment—but this opportunity narrows as structural failure develops.

The literature therefore supports a shift from a purely symptomatic approach toward a **risk-informed, imaging-supported strategy.**

Three dimensions are particularly important:

1. Aetiology

What caused or predisposed to the osteonecrosis?

2. Biology

How extensive is the necrotic lesion?

3. Mechanics

How likely is the lesion to fail under physiological loading?

This third dimension explains why lesion location and size matter so greatly.

The transition from osteonecrosis to collapse is ultimately a biological-mechanical phenomenon.

34. The Integrative Perspective

An integrative model should not mean combining unrelated therapies indiscriminately.

Instead, it should establish a hierarchy of clinical responsibilities:

First: establish the diagnosis.

Second: identify and modify risk factors.

Third: determine stage and collapse risk.

Fourth: provide appropriate orthopedic management.

Fifth: address pain, function, lifestyle, psychological needs and rehabilitation.

Sixth: where desired and clinically appropriate, complementary approaches may be explored with transparent communication regarding the strength and limitations of evidence.

Within this framework, homoeopathic individualization can function as a **patient-centred complementary methodology**, particularly in documenting individualized symptoms and therapeutic preferences, but it should not replace disease surveillance.

35. Clinical Implications for Early Recognition

For clinicians working in primary care, general practice, homoeopathy, physiotherapy or other first-contact settings, the following principle is particularly important:

Persistent unexplained hip or groin pain in a young adult should trigger consideration of osteonecrosis when recognized risk factors are present, even when initial radiographs are unrevealing.

A history of:

- corticosteroid exposure,
- substantial alcohol consumption,
- autoimmune disease,
- hematological disease,
- previous hip trauma,
- transplantation,
- or unexplained bilateral hip symptoms

should increase clinical suspicion.

MRI should be considered when clinical suspicion remains despite nondiagnostic radiography.

36. CONCLUSION

Avascular necrosis of the femoral head is not simply a disorder of “dead bone.” It is the final common pathway of interacting vascular, metabolic, cellular, inflammatory and mechanical disturbances.

Its significance in early adulthood arises from the possibility that an initially silent process can progress toward irreversible structural failure during a period of maximal personal and occupational activity.

The central clinical sequence may be summarized as:

Risk factor → **vascular/microcirculatory injury** → **osteocyte death** → **clinically silent osteonecrosis** → **pain** → **subchondral fracture** → **femoral-head collapse** → **secondary arthritis** → **possible arthroplasty**.

The most important opportunity for intervention lies before irreversible collapse. Early clinical suspicion, MRI-based diagnosis, accurate staging, characterization of lesion size and location, and appropriate orthopedic referral therefore remain fundamental.

Homoeopathic individualization offers a distinct conceptual approach centered on the patient's characteristic symptom pattern, modalities, concomitants and constitutional features. Classical sources emphasize the totality of symptoms and individualized remedy selection. However, there is currently insufficient high-quality evidence to conclude that individualized homoeopathic treatment reverses femoral-head osteonecrosis or prevents structural collapse.

Accordingly, the academically defensible role of homoeopathy in this condition is best framed as complementary and individualized supportive care within an objectively monitored integrative model, rather than as a substitute for evidence-based orthopedic diagnosis and management.

Ultimately, the clinical objective should be not merely to treat pain, but to recognize ischemia before collapse, preserve the native hip whenever realistically possible, reduce modifiable risks, and protect long-term function.

REFERENCES

- Krakowiak J, et al. Osteonecrosis related to steroid and alcohol use—an update on pathogenesis. *Int J Mol Sci*. 2023; 24:11786.
- Zhu W, Guo Z, et al. Treatment of non-traumatic avascular necrosis of the femoral head. *Mol Med Rep*. 2022;25.
- Shah KN, Racine J, Jones LC, Aaron RK. Pathophysiology and risk factors for osteonecrosis. *Curr Rev Musculoskelet Med*. 2015; 8:201-209.
- Mont MA, Cherian JJ, Sierra RJ, Jones LC, Lieberman JR. Nontraumatic osteonecrosis of the femoral head: where do we stand today? A ten-year update. *J Bone Joint Surg Am*. 2015; 97:1604-1627.
- Zhao D, Xu D, Liu D, Wang B. Relationship between the prevalence of osteonecrosis of the femoral head and steroid use. *Clin Orthop Relat Res*. 2007.
- Fukushima W, Fujioka M, Kubo T, Tamakoshi A, Nagai M, Hirota Y. Nationwide epidemiologic survey of idiopathic osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 2010; 468:2715-2724.
- Mont MA, Zywielski MG, Marker DR, McGrath MS, Ulrich SD. The natural history of untreated asymptomatic osteonecrosis of the femoral head. *J Bone Joint Surg Am*. 2011; 93:2165-2170.
- Ohzono K, Saito M, Sugano N, Takaoka K, Ono K. The fate of nontraumatic avascular necrosis of the femoral head. *Clin Orthop Relat Res*. 1991.
- Steinberg ME, Bands A, Parry S, Hoffman E, et al. The prevalence and natural history of early osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 1995.
- Kerachian MA, Séguin C, Harvey EJ. Glucocorticoids in osteonecrosis of the femoral head: a new understanding of the mechanisms of action. *J Steroid Biochem Mol Biol*. 2009; 114:121-128.
- Kopecky KK, Braunstein EM, Brandt KD, Filo RS, Sledge CB. Apparent avascular necrosis of the hip: appearance and significance of MR findings. *Radiology*. 1991.
- Mitchell DG, Kressel HY, Arger PH, et al. Avascular necrosis of the femoral head: morphologic assessment by MR imaging. *Radiology*. 1987.
- Koo KH, Kim R, Ko GH, et al. Preventing collapse in early osteonecrosis of the femoral head. *J Bone Joint Surg Am*. 1995.
- Nishii T, Sugano N, Ohzono K, Sakai T, Haraguchi K, Yoshikawa H. Progression and cessation of collapse in osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 2002.
- Sugano N, Kubo T, Takaoka K, et al. Diagnostic criteria for non-traumatic osteonecrosis of the femoral head. *Clin Orthop Relat Res*. 1999.
- Zeng H, et al. Accuracy of MRI diagnosis of early osteonecrosis of the femoral head: a meta-analysis and systematic review. *J Orthop Surg Res*. 2018; 13:261.
- Yoon BH, Mont MA, Koo KH, Chen CH, Cheng EY, Cui Q, et al. The 2019 revised version of Association Research Circulation Osseous staging system of osteonecrosis of the femoral head. *J Arthroplasty*. 2020.
- Chinese Orthopaedic Association and international collaborators. Guidelines for clinical diagnosis and treatment of osteonecrosis of the femoral head in adults. *J Orthop Transl*. 2020; 24:103-117.
- Hahnemann S. *Organon of medicine*. 6th ed. 1842. Classical homoeopathic text; principles concerning individualization and totality of symptoms.
- Boericke W. *Pocket manual of homoeopathic materia medica*. 9th ed. Boericke & Runyon; 1927. Classical materia-medica reference including *Calcarea fluorica*, *Calcarea carbonica* and *Fluoricum acidum*.

Creative Commons (CC) License

This article is an open-access article distributed under the terms and conditions of the Creative Commons Attribution–Non-Commercial–No Derivatives 4.0 International (CC BY-NC-ND 4.0) license. This license permits sharing and redistribution of the article in any medium or format for non-commercial purposes only, provided that appropriate credit is given to the original author(s) and source. No modifications, adaptations, or derivative works are permitted under this license.

About the author



Dr. Pratibha Jagannath Pagar is a BHMS and MD in Homoeopathy (Practice of Medicine) and serves as Head and Professor of the Department of Practice of Medicine at Matoshri Homoeopathy College & Hospital, Nashik, Maharashtra, India. She is actively involved in teaching, clinical practice, academic activities, and the training of homoeopathic medical students.